

7 Tips to Stop a Nosebleed Fast - MedicineNet

Posterior nosebleeds can often have external or environmental causes, including: picking or scratching your nose a lot; blowing your nose too hard or too often

Posterior Nosebleed: Causes and Treatments

There is so much bleeding that it is hard to breathe. The person is taking blood thinners, such as warfarin (Coumadin), dabigatran (Pradaxa), rivaroxaban (Xarelto), fondaparinux sodium (Arixtra),...

First Aid Treatment for Nosebleeds - WebMD

Nosebleeds can sometimes be a side effect of medication. Certain medications stop blood clotting, which may cause frequent nosebleeds. These drugs can include blood thinners, such as coumadin or...

Daily nosebleeds: Causes and treatment

Nosebleeds are more likely to occur at high altitudes where the air is drier. Very dry environments

can cause nosebleeds. These conditions can cause the inner lining of the nose to dry out and...

Nosebleeds with clots: Causes and treatments

The most common cause of nosebleeds is dry air. The dryness causes crusting inside the nose, which may itch or become irritated. If your nose is scratched or picked, it can bleed. Other common causes include:

Nosebleed: Symptoms, Causes and Treatment - HTQ

The nosebleed follows an accident, a fall or an injury to your head, including a punch in the face that may have broken your nose When to contact your doctor You experience frequent nosebleeds.

Nosebleeds: First aid - Mayo Clinic

Nosebleeds that need medical attention can come from deeper inside the nose and usually affect adults. They can be caused by: an injury or broken nose. high blood pressure. conditions that affect the blood vessels or how the blood clots. certain medicines, like warfarin. Sometimes the cause of a nosebleed is unknown.